


# **EXHIBIT F**

## Accepted Manuscript



### Host Response to Synthetic Mesh in Women with Mesh Complications

Alexis L. Nolfi, BS, Bryan N. Brown, PhD, Rui Liang, MD, Stacy L. Palcsey, BS,  
Michael J. Bonidie, MD, Steven D. Abramowitch, PhD, Pamela A. Moalli, MD/PhD

PII: S0002-9378(16)30049-7

DOI: 10.1016/j.ajog.2016.04.008

Reference: YMOB 11039

To appear in: *American Journal of Obstetrics and Gynecology*

Received Date: 4 January 2016

Revised Date: 31 March 2016

Accepted Date: 8 April 2016

Please cite this article as: Nolfi AL, Brown BN, Liang R, Palcsey SL, Bonidie MJ, Abramowitch SD, Moalli PA, Host Response to Synthetic Mesh in Women with Mesh Complications, *American Journal of Obstetrics and Gynecology* (2016), doi: 10.1016/j.ajog.2016.04.008.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Submit to AJOG

## 1 HOST RESPONSE TO SYNTHETIC MESH IN WOMEN WITH MESH COMPLICATIONS

2 Alexis L. NOLFI BS<sup>1,2,4</sup>, Bryan N. BROWN PhD<sup>1,2,4</sup>, Rui LIANG MD<sup>1,3</sup>, Stacy L. PALCSEY  
 3 BS<sup>1</sup>, Michael J. BONIDIE MD<sup>3</sup>, Steven D. ABRAMOWITCH PhD<sup>2</sup>, Pamela A. MOALLI  
 4 MD/PhD<sup>1,2,3</sup>

5 1. Magee-Womens Research Institute, Pittsburgh, PA

6 2. Department of Bioengineering, University of Pittsburgh, Pittsburgh, PA

7 3. Department of Obstetrics, Gynecology, and Reproductive Sciences, School of Medicine,  
 8 University of Pittsburgh, Pittsburgh, PA

9 4. McGowan Institute for Regenerative Medicine, University of Pittsburgh, Pittsburgh, PA

## 10 DISCLOSURE

11 Moalli: cooperative research agreement ACell

12 Abramowitch: cooperative research agreement ACell

13 Other authors report no conflict of interest

14  
 15 This work was supported by National Institutes of Health awards R01 HD061811 and  
 16 K12HD043441. The content is solely the responsibility of the authors and does not necessarily  
 17 represent the official views of the National Institutes of Health. The funding source had no  
 18 involvement in the study design, collection of data, analysis of data, interpretation of data,  
 19 writing of the report, or the decision to submit for publication.

20 These findings were presented at the American Urogynecologic Society (AUGS) /International  
 21 Urogynecological Association (IUGA) Joint Scientific Meeting, Washington, DC, July 22-26,  
 22 2014 and the 36th Annual Meeting of American Urogynecologic Society PFD Week, Seattle,  
 23 WA, October 13-17, 2015.

24 Corresponding author:

25 Pamela M. Moalli

26 Magee Women Research Institute

27 Department of Obstetrics and Gynecology

28 204 Craft Ave.

29 Pittsburgh, PA 15219

30 Phone: 412-641- 6052

31 Fax: 412-641-5290

32 Email: [pmoalli@mail.magee.edu](mailto:pmoalli@mail.magee.edu)

33 Word Count

34 Abstract: 444; main text: 2912

ACCEPTED MANUSCRIPT

35 CONDENSATION

36 A persistent macrophage response was observed in mesh tissue complexes excised from women  
37 with mesh complications months to years following implantation, with differences observed in  
38 biochemical markers for exposure vs pain.

39 SHORT VERSION OF TITLE

40 Vaginal macrophage profile in women with mesh complications.

41

ACCEPTED MANUSCRIPT

## 42 ABSTRACT

43 Background: In spite of good anatomical and functional outcomes, urogynecologic  
44 polypropylene meshes used to treat pelvic organ prolapse and stress urinary incontinence are  
45 associated with significant complications, most commonly mesh exposure and pain. Few studies  
46 have been performed that specifically focus on the host response to urogynecologic meshes. The  
47 macrophage has long been known to be the key cell type mediating the foreign body response.

48 Conceptually, macrophages responding to a foreign body can be broadly dichotomized  
49 into M1 pro-inflammatory and M2 pro-remodeling subtypes. A prolonged M1 response is  
50 thought to result in chronic inflammation and the formation of foreign body giant cells with  
51 potential for ongoing tissue damage and destruction. While a limited M2 predominant response  
52 is favorable for tissue integration and ingrowth, excessive M2 activity can lead to accelerated  
53 fibrillar matrix deposition, resulting in fibrosis and encapsulation of the mesh.

54 Objectives: To define and compare the macrophage response in patients undergoing a mesh  
55 excision surgery for the indication of pain versus a mesh exposure.

56 Study design: Patients scheduled to undergo a surgical excision of mesh for pain or exposure at  
57 Magee-Womens Hospital were offered enrollment. Twenty-seven mesh-vagina complexes  
58 removed for the primary complaint of a mesh exposure (N=15) vs pain in the absence of an  
59 exposure (n=12) were compared to 30 full thickness vaginal biopsies from women undergoing  
60 benign gynecologic surgery without mesh. Macrophage M1 pro-inflammatory versus M2 pro-  
61 remodeling phenotypes were examined via immunofluorescent labeling for cell surface markers  
62 CD86 (M1) vs CD206 (M2) and M1 vs M2 cytokines via enzyme-linked immunosorbent assay  
63 (ELISA). The amount of matrix metalloproteinase-2 (MMP-2) and matrix metalloproteinase-9

ACCEPTED MANUSCRIPT

64 (MMP-9) proteolytic enzymes were quantified by zymography and substrate degradation assays,  
65 as an indication of tissue matrix degradation. Statistics were performed using one-way ANOVA  
66 with appropriate post hoc tests, t-tests, and Fisher's Exact test.

67 Results: Twenty-seven mesh-vaginal tissue complexes were excised from 27 different women  
68 with mesh complications - 15 incontinence midurethral slings and 12 prolapse meshes. Upon  
69 histological examination, macrophages surrounded each mesh fiber in both groups, with  
70 predominance of the M1 subtype. M1 and M2 cytokines/chemokines, MMP-9 (pro and active),  
71 and MMP-2 (active) were significantly increased in mesh-vagina explants as compared to vagina  
72 without mesh. Mesh explants removed for exposure had 88.4% higher pro-MMP-9 ( $p=0.035$ )  
73 than those removed for pain. A positive correlation was observed between the profibrotic  
74 cytokine IL-10 and the percentage of M2 cells ( $r=0.697$ ,  $p=0.037$ ) in the pain group.

75 Conclusion: In women with complications, mesh induces a pro-inflammatory response that  
76 persists years after implantation. The increase in MMP-9 in mesh explants removed for exposure  
77 indicates degradation while the positive association between IL-10 and M2 macrophages in mesh  
78 explants removed for pain is consistent with fibrosis.

79

80

81 Keywords: cytokines, inflammatory response, macrophage phenotype, polypropylene mesh

## 82 INTRODUCTION

83 Over the past decade, lightweight, wide-pore polypropylene mesh has been increasingly  
84 used in the repair of pelvic organ prolapse and stress urinary incontinence. In spite of favorable  
85 anatomical and functional outcomes, mesh use has been associated with complications, most  
86 commonly mesh exposure through the vaginal epithelium and pain.<sup>1</sup> Studies of similar meshes  
87 used in hernia repair have demonstrated that all polypropylene meshes induce a prolonged  
88 inflammatory response at the site of implantation.<sup>2-3</sup> The magnitude and type of response is  
89 associated with the development of complications.<sup>3-4</sup> Although it is tempting to extrapolate these  
90 findings to meshes applied to the vagina, data suggests that the host response at these two sites is  
91 distinct.<sup>5</sup>

92 The host response following the placement of any foreign material into the body has been  
93 well described. Following the initial recruitment of neutrophils, macrophages become the  
94 primary immune cell involved in the clearance of debris and the initiation of the host response.<sup>6-7</sup>  
95 While this is an essential initial component of healing, the long term presence of activated  
96 inflammatory cells, such as macrophages at the mesh tissue interface, can negatively impact the  
97 ability of the mesh to function as intended.

98 Macrophages have been classified as having diverse and plastic phenotypes along a  
99 continuum between M1 (classically activated; pro-inflammatory) and M2 (alternatively  
100 activated; remodeling, homeostatic) extremes.<sup>8-10</sup> M1 macrophages are characterized by the  
101 secretion of reactive oxygen species, pro-inflammatory cytokines and chemokines, and can be  
102 identified via the cell surface marker CD86. Persistence of M1 macrophages can lead to tissue  
103 damage and destruction. In contrast, M2 macrophages secrete growth factors and anti-  
104 inflammatory immune modulators and can be identified by the cell surface marker CD206. M2

ACCEPTED MANUSCRIPT

macrophages participate in the constructive healing and remodeling phase of the foreign body response resulting in tissue deposition and in growth.<sup>8, 10-12</sup> However, an overzealous M2 response can also lead to excess tissue deposition and fibrosis.<sup>13</sup> As such, macrophage polarization and plasticity play an important and determinant role in tissue remodeling following injury and the integration of biomaterials.<sup>13-14</sup> Limited data exists on the macrophage response following implantation of urogynecologic meshes<sup>15-16</sup> particularly in regard to implantation in the vagina.

The objective of the current study was to characterize the macrophage response present in patients undergoing mesh excision surgery and to define differences in this response according to the two most common complications - mesh exposure and pain.<sup>1</sup> Mesh-tissue constructs from women undergoing mesh excision surgery for the indication of pain and exposure were compared to full thickness vaginal biopsies from women undergoing prolapse repairs using morphological, biochemical, and immunological endpoints.

## **MATERIALS AND METHODS**

### **Patient Acquisition**

Patients scheduled to undergo surgical excision of mesh as part of a larger study (Magee Mesh Biorepository IRB# 10090194) were offered enrollment. For inclusion in the current study, mesh had to be removed from the anterior or apical compartment for the primary indication of exposure or pain. Mesh exposure was defined as at least 2 mm of mesh visible through the vaginal epithelium and pain was defined as mesh being removed for the primary complaint of pain (with palpation, ambulation, or intercourse) without evidence of exposure. Patients were excluded from the study if presenting with acute infection (fever, worsening pain,



ACCEPTED MANUSCRIPT

128 and pus in area of mesh) or erosion into the bowel or bladder. Patients were also excluded if they  
129 were unable to provide informed consent, on chronic immunosuppressive therapy, or had an  
130 autoimmune disorder. After consent was obtained, baseline demographic data were abstracted  
131 from the electronic medical record including age, race/ethnicity, body mass index (BMI),  
132 gravidity, parity, hormone use, menopausal status, and smoking status (Table 1). Menopausal  
133 status was defined as: pre-menopausal (regular menstrual periods within the last 12 months) and  
134 post-menopausal (no menstrual periods within the last 12 months). Hormone use was defined as  
135 current use of systemic estrogen with or without progesterone or vaginal estrogen for  $\geq$  to 3  
136 months. Smoking was defined as current smoker (yes/no). Operative reports from the initial  
137 mesh surgery were reviewed and the type of mesh recorded

138 On the day of surgery, the excised mesh-tissue complex was placed in a sterile specimen  
139 container, immediately placed on ice, and sent for analysis. Samples from mesh patients were  
140 age, BMI, and menopausal status-matched to full thickness vaginal biopsies obtained from the  
141 anterior vagina at the vaginal apex in mesh-naïve women with stage II or III prolapse with and  
142 without incontinence undergoing pelvic surgery, as described previously (IRB # 0412054).<sup>17</sup>  
143 This group of vaginal biopsies was selected as a control for the present study as there is evidence  
144 that the vaginal biopsies from women with prolapse and incontinence have increased MMPs  
145 relative to women with normal pelvic organ support.<sup>18-19</sup>

#### 146 **Tissue Extract Acquisition and Histological Preparation**

147 Excised tissue-mesh complexes and non-mesh control tissues were extracted in high salt  
148 extraction buffer as described previously.<sup>20</sup> Additional pieces of vagina-mesh-complexes and

ACCEPTED MANUSCRIPT

149 non-mesh control tissue were embedded into O.C.T. compound (Tissue-Tek®, Sakura Finetek  
150 USA, Inc., Torrance, CA), flash frozen in liquid nitrogen, sectioned (7µm) and stored at -80°C.

#### 151 **Cytokine and Chemokine Determination**

152 Quantification of cytokines IL-10, IL-4, TNF-α, IL-12p70, and IL-12p40p70 and  
153 chemokines CXCL10 and CCL17 was performed using commercially available ELISA kits (Life  
154 Technologies, Carlsbad, CA and R&D Systems, Minneapolis, MN, respectively). All samples  
155 were run in duplicate or triplicate using 40ug total protein per sample per assay. A patient sample  
156 that had been previously characterized for analyte amounts served as an internal control.

#### 157 **Zymographic Analyses**

158 Samples containing 30µg total protein were analyzed in duplicate by substrate  
159 zymography according to manufacturer's instructions (Novex, Life Technologies, Carlsbad,  
160 CA). For quantification of active and pro-enzyme forms of MMP-2, band density was measured  
161 with ImageJ Software (National Institute of Health, Bethesda, MD), and measurements were  
162 normalized to an internal control.

#### 163 **MMP-9 activity assay**

164 Endogenous MMP-9 activity was measured using a Fluorokine E Human Active MMP-9  
165 kit (R&D systems, Minneapolis, MN). For each experiment, 20 µg of protein from each sample  
166 was tested in duplicate. Fluorescence was read using a Spectramax M2 spectrophotometer  
167 (Molecular Devices, Sunnyvale, CA) at an excitation wavelength of 320 nm and an emission  
168 wavelength of 405 nm. Data was analyzed using a four-parameter regression curve (Masterplex  
169 ReaderFit, Miraibio, San Francisco, CA) and normalized to nanogram active MMP-9 per mg  
170 protein in each sample.

ACCEPTED MANUSCRIPT

**171 Immunofluorescent Labeling**

172 Tissue sections were quadruple-labeled for pan-macrophage marker CD68, M1  
173 macrophage marker CD86, M2 macrophage marker CD206, and nuclear marker 4',6-diamidino-  
174 2-phenylindole (DAPI) as described<sup>15</sup> and imaged using a Nikon ECLIPSE 90i upright  
175 microscope (Nikon USA, Melville, NY). For mesh-vagina complexes, six 200X images were  
176 acquired over two locations within the tissue. Three images were taken in an area of a mesh fiber  
177 (defined as a single fiber of polypropylene not immediately adjacent to other fibers) and three  
178 additional images were taken in the area of a mesh knot (defined as three or more single fibers  
179 immediately adjacent to each other). For control patients, six 200X images were taken over two  
180 areas of each tissue. For each image, two trained technicians counted the number of total cells  
181 and the number of cells co-expressing either CD68 and CD86 or CD68 and CD206 to define the  
182 M1 and M2 macrophage population, respectively. For each field the M2:M1 ratio was calculated  
183 as described in Wolf et al<sup>21</sup> by using the formula (raw number of M2 macrophages+1)/(raw  
184 number of M1 macrophages +1) to avoid division by 0 in samples with no cells present.

**185 Statistical Analysis**

186 Power analysis showed 8 samples in each group were necessary to reach statistical  
187 significance for cytokines using previously obtained IL-10 values in vaginal extracts from non-  
188 human primate with and without Gynemesh PS implanted via sacrocolpopexy.<sup>15</sup> Statistical  
189 analysis was performed with SPSS 21 (IBM, Armonk, NY). For demographic data, a one-way  
190 ANOVA with Tukey *post hoc* testing, Kolmogorov-Smirnov tests, and Fisher's Exact tests with  
191 significance level  $\alpha=0.05$  was used. For the biochemical endpoints of mesh patients vs. control  
192 patients and for the complication exposure vs. pain, a two-tailed Student's independent samples  
193 t-test was performed. For histological endpoints comparing areas of a mesh fiber to a mesh knot,

ACCEPTED MANUSCRIPT

194 a paired-sample t-test was used ( $\alpha=0.05$ ), and when comparing tissue removed for exposure vs.  
195 pain, an independent-sample t-test was used ( $\alpha=0.05$ ). Pearson's correlations were used to  
196 correlate cytokines with histologic findings.

197

## 198 RESULTS

### 199 Demographic Data

200 Twenty-seven mesh-vaginal tissue complexes were excised from 27 women - 15  
201 incontinence midurethral slings and 12 prolapse meshes. Four of the 27 meshes were implanted  
202 via abdominal sacrocolpopexy and the remainder were inserted transvaginally. There were no  
203 differences in patient age, race/ethnicity, BMI, gravidity, parity, hormone use, menopausal  
204 status, smoking status, or duration of mesh implantation (all  $p>0.05$ , Table 1). In addition, there  
205 were no differences in patient demographics or experimental endpoints when separated by  
206 prolapse meshes vs midurethral slings. The specific meshes that were excised are listed in Table  
207 2. Mesh type was not found in 3 meshes removed for pain and 1 mesh removed for exposure.  
208 Meshes were explanted from 4.5 to 93 months following the index surgery.

### 209 Biochemical Endpoints

210 All M1 pro-inflammatory and M2 pro-remodeling cytokines and chemokines were  
211 increased in mesh explants as compared to non-mesh tissue (Table 3), indicating a robust, active  
212 and ongoing host response to polypropylene long after implantation. Examination of the pro-  
213 remodeling cytokines IL-4 and IL-10 showed a 1.18 fold increase ( $p=0.011$ ) and 1.45 fold  
214 increase ( $p=0.016$ ), respectively, in mesh-vagina explants vs. control. Pro-inflammatory  
215 cytokines TNF- $\alpha$ , IL-12p40p70, and IL-12p70 had a 2.13 fold increase ( $p<0.001$ ), a 2.19 fold  
216 increase ( $p<0.001$ ), and a 1.22 fold increase ( $p=0.001$ ), respectively, in mesh explants vs.

ACCEPTED MANUSCRIPT

217 controls. The pro-inflammatory chemokine CXCL10 and the pro-remodeling chemokine CCL17  
 218 were 3.38 fold increased ( $p=0.016$ ) and 7.26 fold increased ( $p<0.001$ ) relative to control,  
 219 respectively. Comparison of the ratio of the M2 pro-remodeling cytokines (IL-10 + IL-4) to the  
 220 M1 pro-inflammatory cytokines (TNF- $\alpha$  + IL-12p70) revealed a decrease in mesh explants as  
 221 compared to controls ( $p=0.003$ ), indicating a shift towards a pro-inflammatory profile.

222 Pro- and active MMP-9 were 2.85 fold increased ( $p<0.0001$ ) and 2.91 fold increased  
 223 ( $p<0.0001$ ) respectively, in mesh explants as compared to controls. While proMMP-2 was  
 224 similar in complexes with and without mesh, active MMP-2 was 2.08 fold increased ( $p=0.038$ ).

225 Pro-MMP-9 was 1.88 fold higher with exposure than pain ( $p=0.036$ ). No statistical  
 226 difference was observed for active MMP-9 ( $p=0.067$ ). Values for individual cytokines and  
 227 chemokines and for levels of pro- and active MMP2 did not differ based on indication for mesh  
 228 removal (Table 4).

## 229 Immunofluorescent Labeling

230 Mesh tissue complexes demonstrated a marked but highly localized foreign body  
 231 response characterized by the presence of CD68+ cells (macrophages) surrounding each mesh  
 232 fiber. In areas where mesh fibers were in close proximity, the host response to neighboring fibers  
 233 overlapped increasing the magnitude of the response. Labeling with the M1-specific surface  
 234 marker CD86 and the M2-specific marker CD206 demonstrated that control tissues contained  
 235 few or no macrophages, as opposed to tissue-mesh-complexes in which CD68+CD86+ pro-  
 236 inflammatory M1 macrophages were concentrated around mesh fibers (Figure 1, Table 5). Mesh  
 237 explants contained a higher number of total cells/200X field when compared to controls (682.46

ACCEPTED MANUSCRIPT

238  $\pm 142.61$  vs  $441.63 \pm 126.13$  cells,  $p < 0.001$ ) and a lower ratio of M2:M1 macrophages ( $0.260 \pm$   
239  $0.161$  vs  $1.772 \pm 1.919$ ,  $p = 0.001$ ), supporting an ongoing pro-inflammatory response.

240 In mesh explants, no differences in the number of macrophages present or the  
241 macrophage phenotype was observed when data was stratified based on indication for mesh  
242 removal (Table 6). However, when comparing areas containing a mesh fiber vs. a mesh knot in  
243 individual samples, the area of a mesh knot had a total cell density that was 1.23 fold increased  
244 ( $p = 0.037$ ). Within the area of a mesh knot, the number of M1 macrophages was 2.00 fold  
245 increased ( $p = 0.003$ ), the percentage of M1 macrophages (as a function of total cells) was 1.59  
246 fold increased ( $p = 0.004$ ), the number of M2 cells was 2.27 fold increased ( $p < 0.001$ ), and  
247 percentage of M2 macrophages was 1.77 fold increased ( $p = 0.002$ ) as compared to a single mesh  
248 fiber (Table 7) suggesting that the host response was proportional to the amount of material in  
249 contact with the host. No significant difference was observed in the ratio of M2:M1  
250 macrophages in the area of a mesh knot vs a mesh fiber. A positive correlation between IL-10  
251 and the percentage of pro-remodeling/pro-fibrotic M2 cells ( $r = 0.697$ ,  $p = 0.037$ ) in the pain group  
252 was observed that was not present in the exposure group. In the exposure group, a positive  
253 correlation was observed between the pro-inflammatory cytokine IL-12p40p70 and percentage of  
254 pro-inflammatory M1 macrophages ( $r = 0.584$ ,  $p = 0.059$ ), which did not reach statistical  
255 significance. This correlation was not observed in the pain group.

## 256 COMMENT

257 A persistent foreign body response was observed in mesh tissue complexes excised from  
258 women requiring surgical excision of mesh months to years after mesh implantation. The host  
259 response was characterized by a predominance of macrophages with an increase in both pro-

ACCEPTED MANUSCRIPT

260 inflammatory and pro-remodeling cytokines/chemokines along with increased tissue degradation  
261 as evidenced by increased MMP-2 and MMP-9. Mesh-tissue complexes removed for mesh  
262 exposure had increased pro-MMP-9 indicating a pro-inflammatory and tissue destruction type  
263 response. In contrast, in mesh-tissue complexes removed for pain, the percentage of M2  
264 macrophages (involved in tissue remodeling and fibrosis<sup>22-24</sup>) positively correlated with the  
265 amount of the anti-inflammatory/pro-fibrotic cytokine IL-10 consistent with tissue deposition  
266 and encapsulation.

267 The presence of macrophages, elevated cytokines, chemokines and MMPs in tissue-mesh  
268 complexes excised from patients with exposure or pain suggests that polypropylene mesh elicits  
269 an ongoing host inflammatory response.<sup>7</sup> Importantly, the presence of macrophages was limited  
270 to the area immediately surrounding the mesh fibers with each fiber eliciting an independent  
271 reaction, the magnitude of which appeared to be proportional to the number of fibers in a given  
272 area. This points to the importance of maintaining meshes in a flat (as opposed to folded)  
273 configuration to minimize the amount of material per area and choosing meshes in which the  
274 spaces between fibers (pores) are wide enough that the host response to two adjacent fibers does  
275 not overlap.<sup>25</sup>

276 In mesh-tissue complexes removed for exposure, pro-MMP9 was increased as compared  
277 to mesh removed for pain, suggesting a degenerative process. Studies of hernia meshes in animal  
278 models have shown that persistent inflammation with a prolonged release/activation of MMPs  
279 can lead to the degeneration of mesh-implanted tissues resulting in a deterioration in structural  
280 and mechanical integrity.<sup>26-27</sup> In a nonhuman primate model, Gynemesh PS caused a decrease in  
281 key structural proteins (collagen and elastin) and increased MMP activity leading to thinning of  
282 the underlying and associated tissues and a deterioration of mechanical properties.<sup>20, 28-29</sup>



ACCEPTED MANUSCRIPT

283 Gynemesh PS has a highly unstable geometry when loaded resulting in pore collapse and  
284 increasing stiffness of the product.<sup>30-31</sup> As virtually all meshes removed from women with  
285 complications have evidence of deformation and pore collapse,<sup>32-36</sup> mesh exposure may  
286 represent a mechanical phenomenon in which altered mesh geometries result in increased mesh  
287 stiffness which in turn, leads to a degenerative response as a result of stress shielding and  
288 ongoing destructive inflammation.

289 While the mechanism for developing pain following mesh implantation is not clear, mesh  
290 deformation (contraction, retraction, or shrinkage) is also frequently observed in meshes  
291 removed for pain.<sup>32, 34, 36-38</sup> In normal wound healing, as inflammation resolves and remodeling  
292 begins, some amount of tissue contraction occurs with the formation of a scar.<sup>38</sup> In the presence  
293 of a permanent foreign body, the implant is surrounded with a fibrotic capsule since it cannot be  
294 degraded. For hernia meshes, if the fibers are too close (<1mm), the fibrotic response to  
295 neighboring fibers overlaps, or “bridges,” resulting in “bridging fibrosis” or encapsulation of the  
296 mesh.<sup>25</sup> As myofibroblasts constitute the primary cellular component of the fibrotic capsule,  
297 when a mesh becomes encapsulated, the resulting contraction or “shrinkage” may place tension  
298 on adjacent tissues resulting in pain. Indeed, mesh shrinkage (50-70%) has been described to  
299 occur following transvaginal insertion of prolapse meshes.<sup>32-37</sup> In the present study, in meshes  
300 removed for pain, IL-10, a cytokine that, in increased amounts, has been associated with  
301 fibrosis,<sup>39-42</sup> positively correlated with the percentage of M2 polarized macrophages  
302 (remodeling/fibrotic phenotype), supporting an ongoing remodeling/fibrotic process involved in  
303 at least one mechanism leading to pain.

304 A major limitation of the current study is that it was not possible to include a control group  
305 of mesh-tissue complexes obtained from women who underwent mesh implantation without a



ACCEPTED MANUSCRIPT

306 complication. As such, the current study does not assess the inflammatory response to prolapse  
307 mesh in women with a good outcome and focuses only upon the inflammatory response in the  
308 setting of complications. Our strict inclusion criteria resulted in a limited sample size which  
309 limits the generalizability of the data especially in regards to the impact of variables such as age,  
310 length of time of mesh implantation, and hormone use. All patients enrolled in the current study  
311 completed the Pelvic Pain Scale (PPS) and a number of other validated questionnaires, the  
312 analysis of which is pending and will be the focus of a future manuscript. Finally, it is our  
313 practice to remove as much mesh as possible in patients presenting with a mesh complication;  
314 therefore, we cannot guarantee that the mesh undergoing analysis was at the exact site of the  
315 complication, regardless of whether it was removed for an exposure or pain.

316 In conclusion, the findings of present study suggest that the two major mesh  
317 complications - exposure and pain - are associated with a marked pro-inflammatory response that  
318 persists years after mesh implantation. In addition, different mechanisms for mesh exposure and  
319 pain may be associated with differential macrophage activation. Future studies will focus on the  
320 specific risk factors that predispose to specific types of mesh complications.

#### 321 **AUTHOR CONTRIBUTIONS**

322 A.L.N. acquiring samples, conducting experiments, writing the manuscript, and analyzing data;  
323 B.B. analyzing data and writing the manuscript; R.L. acquiring data, analyzing data, and writing  
324 the manuscript; S.P. acquiring samples, and conducting experiments; M.B. acquiring data and  
325 analyzing data; S.A. acquiring data and analyzing data; P.A.M. designing research studies,  
326 analyzing data, and writing the manuscript.

#### 327 **ACKNOWLEDGEMENT**

ACCEPTED MANUSCRIPT

328 Funding from R01 HD061811 (P.A.M) and K12 HD043441 (B.N.B)

329

# 330 REFERENCES

- 331 1. FS C. UPDATE on Serious Complications Associated with Transvaginal Placement of  
332 Surgical Mesh for Pelvic Organ Prolapse. In: FDA, editor. July 2011.
- 333 2. Cervigni M, Natale F. The use of synthetics in the treatment of pelvic organ prolapse.  
334 Curr Opin Urol. 2001;11(4):429-35.
- 335 3. Klinge U, Klosterhalfen B, Muller M, Schumpelick V. Foreign body reaction to meshes  
336 used for the repair of abdominal wall hernias. Eur J Surg. 1999;165(7):665-73.
- 337 4. Cobb WS, Kercher KW, Heniford BT. The argument for lightweight polypropylene mesh  
338 in hernia repair. Surgical innovation. 2005;12(1):63-9.
- 339 5. Pierce LM, Rao A, Baumann SS, Glassberg JE, Kuehl TJ, Muir TW. Long-term  
340 histologic response to synthetic and biologic graft materials implanted in the vagina and  
341 abdomen of a rabbit model. Am J Obstet Gynecol. 2009;200(5):546 e1-8.
- 342 6. Anderson JM. Biological Responses to Materials. Annu Rev Mater Res. 2001;31:81-110.
- 343 7. Anderson JM, Rodriguez A, Chang DT. Foreign body reaction to biomaterials. Semin  
344 Immunol. 2008;20(2):86-100. PMCID: 2327202.
- 345 8. Mantovani A, Sica A, Sozzani S, Allavena P, Vecchi A, Locati M. The chemokine  
346 system in diverse forms of macrophage activation and polarization. Trends Immunol.  
347 2004;25(12):677-86.
- 348 9. Mills CD, Kincaid K, Alt JM, Heilman MJ, Hill AM. M-1/M-2 macrophages and the  
349 Th1/Th2 paradigm. J Immunol. 2000;164(12):6166-73.
- 350 10. Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. Nat Rev  
351 Immunol. 2008;8(12):958-69. PMCID: 2724991.

ACCEPTED MANUSCRIPT

- 352 11. Brown BN, Badylak SF. Expanded applications, shifting paradigms and an improved  
353 understanding of host-biomaterial interactions. *Acta Biomater.* 2013;9(2):4948-55.
- 354 12. Gundra UM, Girgis NM, Ruckerl D, et al. Alternatively activated macrophages derived  
355 from monocytes and tissue macrophages are phenotypically and functionally distinct. *Blood.*  
356 2014;123(20):e110-22. PMCID: 4023427.
- 357 13. Murray PJ, Wynn TA. Protective and pathogenic functions of macrophage subsets. *Nat*  
358 *Rev Immunol.* 2011;11(11):723-37. PMCID: 3422549.
- 359 14. Brown BN, Sicari BM, Badylak SF. Rethinking regenerative medicine: a macrophage-  
360 centered approach. *Front Immunol.* 2014;5:510. PMCID: 4219501.
- 361 15. Brown BN, Mani D, Nolfi AL, Liang R, Abramowitch S, Moalli PA. Characterization of  
362 the host inflammatory response following implantation of prolapse mesh in rhesus macaque. *Am*  
363 *J Obstet Gynecol.* 2015.
- 364 16. Hill AJ, Unger CA, Solomon ER, Brainard JA, Barber MD. Histopathology of excised  
365 midurethral sling mesh. *Int Urogynecol J.* 2015;26(4):591-5.
- 366 17. Zong W, Stein SE, Starcher B, Meyn LA, Moalli PA. Alteration of vaginal elastin  
367 metabolism in women with pelvic organ prolapse. *Obstet Gynecol.* 2010;115(5):953-61.  
368 PMCID: 3042770.
- 369 18. Chen BH, Wen Y, Li H, Polan ML. Collagen metabolism and turnover in women with  
370 stress urinary incontinence and pelvic prolapse. *Int Urogynecol J Pelvic Floor Dysfunct.*  
371 2002;13(2):80-7; discussion 7.
- 372 19. Moalli PA, Shand SH, Zyczynski HM, Gordy SC, Meyn LA. Remodeling of vaginal  
373 connective tissue in patients with prolapse. *Obstet Gynecol.* 2005;106(5 Pt 1):953-63.

ACCEPTED MANUSCRIPT

- 374 20. Liang R, Zong W, Palesey S, Abramowitch S, Moalli PA. Impact of prolapse meshes on  
 375 the metabolism of vaginal extracellular matrix in rhesus macaque. *Am J Obstet Gynecol*.  
 376 2015;212(2):174.e1-7.
- 377 21. Wolf MT, Dearth CL, Ranallo CA, et al. Macrophage polarization in response to ECM  
 378 coated polypropylene mesh. *Biomaterials*. 2014;35(25):6838-49.
- 379 22. Novak ML, Koh TJ. Macrophage phenotypes during tissue repair. *Journal of leukocyte*  
 380 *biology*. 2013;93(6):875-81. PMCID: 3656331.
- 381 23. Venosa A, Malaviya R, Choi H, Gow AJ, Laskin JD, Laskin DL. Characterization of  
 382 Distinct Macrophage Subpopulations During Nitrogen Mustard-Induced Injury and Fibrosis. *Am*  
 383 *J Respir Cell Mol Biol*. 2015.
- 384 24. Wynn TA, Barron L. Macrophages: master regulators of inflammation and fibrosis.  
 385 *Semin Liver Dis*. 2010;30(3):245-57. PMCID: 2924662.
- 386 25. Orenstein SB, Saberski ER, Kreutzer DL, Novitsky YW. Comparative analysis of  
 387 histopathologic effects of synthetic meshes based on material, weight, and pore size in mice. *The*  
 388 *Journal of surgical research*. 2012;176(2):423-9.
- 389 26. Jones JA, McNally AK, Chang DT, et al. Matrix metalloproteinases and their inhibitors  
 390 in the foreign body reaction on biomaterials. *Journal of biomedical materials research Part A*.  
 391 2008;84(1):158-66.
- 392 27. Klinge U, Junge K, Stumpf M, Ap AP, Klosterhalfen B. Functional and morphological  
 393 evaluation of a low-weight, monofilament polypropylene mesh for hernia repair. *J Biomed Mater*  
 394 *Res*. 2002;63(2):129-36.

ACCEPTED MANUSCRIPT

- 395 28. Feola A, Abramowitch S, Jallah Z, et al. Deterioration in biomechanical properties of the  
396 vagina following implantation of a high-stiffness prolapse mesh. BJOG. 2013;120(2):224-32.  
397 PMID: 3530836.
- 398 29. Liang R, Abramowitch S, Knight K, et al. Vaginal degeneration following implantation  
399 of synthetic mesh with increased stiffness. BJOG. 2013;120(2):233-43. PMID: 3531826.
- 400 30. Barone W, Moalli P, Abramowitch S. Variable Porosity of Common Prolapse Mesh's  
401 during Uni-Axial Loading. 34th Annual Meeting of American Urogynecologic Society; October  
402 16-19; Las Vegas, NV2013. p. S56-7.
- 403 31. Otto J, Kaldenhoff E, Kirschner-Hermanns R, Muhl T, Klinge U. Elongation of textile  
404 pelvic floor implants under load is related to complete loss of effective porosity, thereby favoring  
405 incorporation in scar plates. Journal of biomedical materials research Part A. 2014;102(4):1079-  
406 84.
- 407 32. Caquant F, Collinet P, Debodinance P, et al. Safety of Trans Vaginal Mesh procedure:  
408 retrospective study of 684 patients. J Obstet Gynaecol Res. 2008;34(4):449-56.
- 409 33. Rogowski A, Bienkowski P, Tosiak A, Jerzak M, Mierzejewski P, Baranowski W. Mesh  
410 retraction correlates with vaginal pain and overactive bladder symptoms after anterior vaginal  
411 mesh repair. Int Urogynecol J. 2013;24(12):2087-92.
- 412 34. Svabik K, Martan A, Masata J, El-Haddad R, Hubka P, Pavlikova M. Ultrasound  
413 appearances after mesh implantation--evidence of mesh contraction or folding? Int Urogynecol J.  
414 2011;22(5):529-33.
- 415 35. Tunn R, Picot A, Marschke J, Gauruder-Burmester A. Sonomorphological evaluation of  
416 polypropylene mesh implants after vaginal mesh repair in women with cystocele or rectocele.  
417 Ultrasound Obstet Gynecol. 2007;29(4):449-52.

ACCEPTED MANUSCRIPT

- 418 36. Velemir L, Amblard J, Fatton B, Savary D, Jacquetin B. Transvaginal mesh repair of  
419 anterior and posterior vaginal wall prolapse: a clinical and ultrasonographic study. *Ultrasound*  
420 *Obstet Gynecol.* 2010;35(4):474-80.
- 421 37. Eisenberg VH, Steinberg M, Weiner Z, et al. Three-dimensional transperineal ultrasound  
422 for imaging mesh implants following sacrocolpopexy. *Ultrasound Obstet Gynecol.*  
423 2014;43(4):459-65.
- 424 38. Marschke J, Hengst L, Schwertner-Tiepelmann N, Beilecke K, Tunn R. Transvaginal  
425 single-incision mesh reconstruction for recurrent or advanced anterior vaginal wall prolapse.  
426 *Arch Gynecol Obstet.* 2014.
- 427 39. Desmouliere A, Chaponnier C, Gabbiani G. Tissue repair, contraction, and the  
428 myofibroblast. *Wound Repair Regen.* 2005;13(1):7-12.
- 429 40. Gerber JS, Mosser DM. Reversing lipopolysaccharide toxicity by ligating the  
430 macrophage Fc gamma receptors. *J Immunol.* 2001;166(11):6861-8.
- 431 41. Mosser DM. The many faces of macrophage activation. *Journal of leukocyte biology.*  
432 2003;73(2):209-12.
- 433 42. Sziksz E, Pap D, Lippai R, et al. Fibrosis Related Inflammatory Mediators: Role of the  
434 IL-10 Cytokine Family. *Mediators Inflamm.* 2015;2015:764641. PMCID: 4495231.

435

ACCEPTED MANUSCRIPT

436 **Table 1.** Descriptive statistics of study population.

437

	Mesh complications (n=27)			
	Mesh exposure (n=15)	Pain (n=12)	Prolapse (n=30)	P-value
Age <sup>a</sup> (years)	56.1 ± 8.0	52.1 ± 9.7	52.9 ± 9.2	0.527
BMI <sup>a</sup>	30.8 ± 5.9	27.3 ± 3.3	28.5 ± 4.0	0.158
Gravidity <sup>b</sup>	3.0 (2.0, 4.0)	3.5 (2.0, 5.75)	2.0 (2.0, 3.0)	0.644
Parity <sup>b</sup>	2.0 (2.0, 3.0)	3.0 (2.0, 5.0)	2.0 (2.0, 3.0)	0.899
Time implanted (months) <sup>a</sup>	36.9 ± 30.3	30.9 ± 18.0	NA	0.527
Menopausal status <sup>c</sup>				
pre-menopausal	2(13%)	3(25%)	12(40%)	0.190
post-menopausal	13(87%)	9(75%)	18(60%)	
Smoking <sup>c</sup>				
nonsmoker	12(80%)	9(75%)	27(90%)	0.713
smoker	3(20%)	3(25%)	3(10%)	
Race/ethnicity <sup>c</sup>				
white	15(100%)	12(100%)	29(97%)	1.000
Indian	0	0	1(3%)	
other	0	0	0	
Hormonal Usage <sup>c</sup>				
Yes	7(47%)	7(58%)	7(23%)	0.077
No	8(53%)	5(42%)	23(77%)	

438 a. values given as mean ± standard deviation

439 b. values given as median (25 percentile, 75 percentile)

440 c. values given as number of patients (percentage)

441

442

443

444

445

446

ACCEPTED MANUSCRIPT

447 **Table 2.** Excised mesh brand and type categorized by mesh complications. Of the 27 patients  
 448 enrolled in this study, mesh brand information was not able to be determined for 4 of the  
 449 patients.

Mesh device	Removal due to exposure (n=15)	Removal due to pain (n=12)
AMS Monarc TOT	1	0
AMS Perigee	1	0
Bard Ajust Single Incision Sling (SIS)	0	1
Bard Soft Mesh	0	1
Boston Scientific Lynx TVT	1	1
Boston Scientific Obtryx TOT	1	0
Boston Scientific Solyx mini sling	0	1
Boston Scientific Uphold	1	0
Caldera Desara Sling System for SUI	1	0
Coloplast Novasilk	1	0
Gynecare Prolift mesh kit	1	1
Gynecare Gynemesh PS	4	0
Gynecare TVT Secur	1	2
Gynecare TVT	1	1
Gynecare TOT	0	1
Original medical records not available	1	3



ACCEPTED MANUSCRIPT

450 **Table 3.** Distribution of Biomarkers by Study Group.

	Mesh complications (exposure + pain) (n=27)	Prolapse control (n=30)	P-Value
<b>Cytokines &amp; Chemokines*</b>			
IL-4	3.316 ± 0.860	2.800 ± 0.567	0.011
IL-10	17.738 ± 9.957	12.218 ± 5.932	0.016
TNF-α	15.963 ± 6.392	7.487 ± 2.224	0.000
IL-12p40p70	28.519 ± 16.437	13.029 ± 7.227	0.000
IL-12p70	2.070 ± 0.471	1.691 ± 0.291	0.001
CXCL10	62.534 ± 71.835	18.492 ± 61.065	0.016
CCL17	33.940 ± 38.156	4.674 ± 3.681	0.000
<b>Proteases</b>			
Pro-MMP-2 <sup>±</sup>	1.192 ± 0.851	1.324 ± 0.971	0.601
Active MMP-2 <sup>±</sup>	1.521 ± 1.636	0.733 ± 0.783	0.038
Pro-MMP-9 <sup>γ</sup>	3.496 ± 2.643	1.226 ± 1.715	0.000
<b>M2/M1 cytokines &amp; chemokines</b>			
(IL-4+IL-10)/ (TNF-α +IL-12p70)	1.209 ± 0.562	1.633 ± 0.486	0.003
CCL17/CXCL10	0.852 ± 0.921	0.661 ± 0.597	0.364

451 Data expressed as Mean ± Standard Deviation

452 \* pg per 40μg protein, <sup>±</sup> arbitrary units, <sup>γ</sup> ng per 15μg total protein

ACCEPTED MANUSCRIPT

453 **Table 4.** Comparison of the profiles of cytokines, chemokines and MMP2 in vaginal tissue  
 454 excised for mesh exposure and pain.

	Mesh: Exposure	Mesh: Pain	P-Value
<b>Cytokines &amp; Chemokines*</b>			
IL-4	3.307 ± 0.973	3.328 ± 0.736	0.952
IL-10	19.323 ± 11.665	15.757 ± 7.308	0.342
TNF- $\alpha$	15.954 ± 5.011	15.976 ± 8.039	0.994
IL-12p40p70	31.739 ± 20.242	24.493 ± 9.235	0.231
IL-12p70	2.040 ± 0.564	2.108 ± 0.342	0.705
CXCL10	71.384 ± 86.873	51.472 ± 48.495	0.459
CCL17	38.445 ± 49.238	28.312 ± 17.132	0.467
<b>Proteases</b>			
Pro-MMP-2 <sup>±</sup>	0.965 ± 1.018	1.420 ± 0.606	0.200
Active MMP-2 <sup>±</sup>	1.278 ± 0.887	1.764 ± 2.164	0.483
Pro-MMP-9 <sup>γ</sup>	4.860 ± 3.464	2.586 ± 1.660	0.036
Active MMP-9 <sup>γ</sup>	6.063 ± 4.674	3.578 ± 1.338	0.067
<b>M2/M1 cytokines &amp; chemokines</b>			
(IL-4+IL-10)/			
	1.266 ± 0.640	1.138 ± 0.462	0.551
(TNF- $\alpha$ +IL-12p70)			
CCL17/CXCL10	0.864 ± 1.165	0.837 ± 0.529	0.937

455 Data expressed as Mean ± Standard Deviation

456 \* pg per 40 $\mu$ g protein, <sup>±</sup> arbitrary units, <sup>γ</sup>ng per 15 $\mu$ g total protein

ACCEPTED MANUSCRIPT

**Table 5.** Macrophage phenotypes in vaginal tissue excised for mesh complications as compared to tissue derived from prolapse women.

	Mesh complications (n=20)	Prolapse (n=24)	P-value
Total Nuclei	682.46 ± 142.61	441.63 ± 126.13	0.000
M1 cell count	197.95 ± 97.15	0.889 ± 2.40	0.000
% M1 cells	28.06 ± 11.03	0.181 ± 0.469	0.000
M2 cell count	41.90 ± 20.33	1.067 ± 2.034	0.000
% M2 cells	6.15 ± 2.80	0.313 ± 0.756	0.000
M2/M1 ratio	0.260 ± 0.161	1.772 ± 1.919	0.001

Data expressed as Mean ± Standard Deviation

ACCEPTED MANUSCRIPT

469 **Table 6.** Comparison of macrophage phenotypic profile in vaginal tissue excised for mesh  
 470 exposure and pain.

	Exposure (n=11)	Pain (n=9)	P-Value
Total Nuclei	660.242 ± 130.892	709.615 ± 159.307	0.467
M1 cell count	167.008 ± 87.598	235.763 ± 99.497	0.124
% M1 cells	24.554 ± 10.669	32.344 ± 10.432	0.118
M2 cell count	40.296 ± 21.046	43.867 ± 20.494	0.706
% M2 cells	6.139 ± 3.157	6.164 ± 2.474	0.984
M2/M1 ratio	0.301 ± 0.206	0.210 ± 0.061	0.189

471 Data expressed as Mean ± Standard Deviation

472

ACCEPTED MANUSCRIPT

473 **Table 7.** Comparison of macrophage phenotypic profile in areas of mesh knots and fibers.

	Mesh knot (n=13)	Mesh fiber (n=13)	P-Value
Total Nuclei	659.87 ± 133.57	536.77 ± 168.30	0.037
M1 cell count	197.79 ± 97.02	98.79 ± 58.79	0.003
% M1 cells	28.89 ± 11.63	18.13 ± 9.88	0.004
M2 cell count	47.08 ± 20.20	20.72 ± 13.24	<0.0001
% M2 cells	7.17 ± 2.83	4.06 ± 2.45	0.002
M2/M1 ratio	0.303 ± 0.186	0.274 ± 0.139	0.715

474

475

476

477

478

ACCEPTED MANUSCRIPT

479 **FIGURE LEGENDS**

480 Figure 1. Immunofluorescent labeling of pan-macrophage marker CD68 (red), M1 pro-  
481 inflammatory marker CD86 (orange), M2 pro-remodeling macrophage marker CD206 (green),  
482 and DAPI (blue). A, a mesh-tissue section from a patient presenting with an exposure and  
483 implanted with the AMS Perigee prolapse mesh for 93 months; B, a mesh-tissue section from a  
484 patient presenting with pain and implanted with the Gynecare TVT Secur for 6 months; C,  
485 control tissue from patients without graft implantation. A predominance of pro-inflammatory M1  
486 macrophages surround mesh fibers (\*) consistent with a prolonged immune response could be  
487 observed in both A and B; however, this response is limited to the area immediately adjacent to  
488 mesh fibers. Control tissue contained few or no macrophages as compared to mesh patient tissue.  
489 Magnification 200X.

490

491

ACCEPTED MANUSCRIPT



ACCEPTED MANUSCRIPT